The Hidden Switches Behind Bacterial Cellulose Production

Aisury T. Lan C. and J. L. Strap

Faculty of Science, Ontario Tech University, Oshawa, ON

Abstract

What if bacteria could help produce the next generation of sustainable materials? Bacterial cellulose is a highly pure biomaterial with applications in medicine, food, and eco-friendly packaging. However, improving its production remains a major challenge. *Komagataeibacter xylinus* is an efficient cellulose-producing bacterium and an ideal model for studying this process. Although cellulose biosynthesis is well understood, its regulatory mechanisms remain largely unknown. This research investigates how gene regulation and bacterial physiology influence cellulose production, aiming to identify factors that could improve yields and support more efficient, sustainable production.

Introduction

Komagataeibacter xylinus is a gram-negative bacterium known for its ability to produce cellulose, making it an organism of interest in biotechnology and bioprocessing (Singhania *et al.* 2021; (Subbiahdoss *et al.*, 2022). Transcription factors are proteins that regulate gene expression by binding to specific DNA sequences, controlling whether genes are activated or repressed. They help bacteria respond to environmental changes by coordinating cellular processes such as metabolism, growth, and stress adaptation (Wang *et al.,* 2015).




Figure 1. Glucose uptake and cellulose biosynthesis pathway in *Komagataeibacter xylinus*. Glucose is transported into the cell and phosphorylated to glucose-6-phosphate, which is then converted into UDP-glucose. The *bcs* operon utilizes UDP-glucose to synthesize bacterial cellulose (BC), which is extruded outside the cell. Adapted from Singhania *et al.* (2021). Originally created in BioRender and subsequently redesigned with assistance from ChatGPT (OpenAI, 2026).




Figure 2. Bacterial cellulose applications. Overview of the different physical forms of bacterial cellulose and its versatility as a biomaterial across diverse industrial, biomedical, and environmental applications. Adapted from Potočnik *et al., (*2023). Originally created using BioRender and subsequently redesigned with assistance from ChatGPT (OpenAI, 2026).

Objective:
To characterize the role of a transcription factor in regulating bacterial physiology, environmental adaptation, and cellulose production in *Komagataeibacter xylinus*.

Methodology




Figure 3. Proposed model of transcription factor-mediated regulation of bacterial cellulose production. Environmental signals, including carbon sources and phytohormones may influence transcription factor activity, potentially modulating the expression of genes involved in BC synthesis. Direct functional validation is required to establish these regulatory relationships.




Figure 4. Construction and selection of *K. xylinus* Δ*gene* mutants. 1-4) The mutator construct (MC) was engineered and introduced into *K. xylinus* electrocompetent cells via electroporation. 5) cells were then recovered in SHG medium and plated on SHG agar supplemented with 50 µg/mL streptomycin and 120 colonies were selected for screening. Integration of the mutator construct into the genome was confirmed by the presence of streptomycin-resistant colonies. This indicated a single crossover event. 6) Colonies were patched twice onto SHG + 5% sucrose plates to select for the second recombination event. Sucrose-resistant colonies indicated excision of the plasmid, resulting in either wild-type or gene-deleted strains. 7-9) Colonies were patched onto SHG plates without antibiotics to confirm plasmid loss and finalize the screening of *K. xylinus Δgene* mutants using PCR and sequencing confirmation. Originally created in BioRender and subsequently redesigned with assistance from ChatGPT (OpenAI, 2026).

Results




Figure 5. Sequencing confirmation of *K. xylinus* Δ*gene* genomic mutants. Genomic DNA from screened colonies suspected to have undergone the double cross-over recombination event were validated by PCR-amplification and sequence analysis. Sequencing reads were aligned to the reference *K. xylinus* ATCC 53582 genome. The genomic region is highlighted in pink. The clean deletion generated by the mutator construct is indicated by the red arrows. Sequence alignments confirm successful recombination at the left and right borders, validating precise excision of the gene at the expected genomic locations.




Figure 6. Preliminary evidence of altered bacterial cellulose production in *K. xylinus*. The Δ*gene* mutant exhibits reduced pellicle formation compared with the wild-type suggesting the deleted gene may contribute to cellulose biosynthesis.




Figure 7. Schematic representation of the proposed regulatory role of the putative regulator in *K. xylinus* and the experimental approaches used to characterize its function. The upper panel illustrates the bacterial cell envelope, quorum-sensing signaling, and the *bcs* operon, which encodes components of the bacterial cellulose biosynthesis machinery. The *bcs* complex facilitates the synthesis and export of cellulose microfibrils into the extracellular matrix. The lower panels illustrate the three main experimental assessments comparing wild-type, Δ*gene* mutant, and complemented strains, A) bacterial cellulose pellicle synthesis, B) antibiotic susceptibility, and C) cell growth kinetics. Originally created in BioRender and subsequently redesigned with assistance from ChatGPT (OpenAI, 2026).

Conclusion

- ✓ The study successfully implemented a clean deletion strategy to remove a regulatory gene in *Komagataeibacter xylinus* using the pKNG101 suicide vector.
- ✓ First functional characterization of this putative regulator in *K. xylinus*.
- ✓ Provides insight into the role of a hybrid regulator in bacterial cellulose-associated processes.

Future Directions

- ➢ Establish whether observed phenotypes are attributable completely to the loss of the putative regulator.
- ➢ Determine the effect of the putative regulator on gene expression and identify associated cellular pathways.
- ➢ Determine whether the putative regulator directly regulates selected candidate target genes.

References

Augimeri, R. V., Varley, A. J., & Strap, J. L. (2015). Establishing a role for bacterial cellulose in environmental interactions: Lessons learned from diverse biofilm-producing Proteobacteria. *Frontiers in Microbiology*, *6*, 1282. https://doi.org/10.3389/fmicb.2015.01282

Deng, Y., Nagachar, N., Xiao, C., Tien, M., & Kao, T. (2013). Identification and characterization of non-cellulose-producing mutants of *Gluconacetobacter hansenii* generated by Tn5 transposon mutagenesis. *Journal of Bacteriology*, *195*(22), 5072–5083. https://doi.org/10.1128/jb.00767-13

Olszakier, S., & Berlin, S. (2022). A simplified Gibson assembly method for site directed mutagenesis by re-use of standard, and entirely complementary, mutagenesis primers. *BMC Biotechnology*, *22*(1), 10. https://doi.org/10.1186/s12896-022-00740-y

Potočnik, V., Gorgieva, S., & Trček, J. (2023). From nature to lab: Sustainable bacterial cellulose production and modification with synthetic biology. *Polymers, 15*(16), 3466. https://doi.org/10.3390/polym15163466

Singhania, R. R., Patel, A. K., Tsai, M.-L., Chen, C.-W., & Dong, C. D. (2021). Genetic modification for enhancing bacterial cellulose production and its applications. *Bioengineered, 12*(1), 6793–6807. https://doi.org/10.1080/21655979.2021.1968989

Subbiahdoss, G., Osmen, S., & Reimhult, E. (2022). Cellulosic biofilm formation of *Komagataeibacter* in kombucha at oil-water interfaces. *Biofilm, 4*, 100071. https://doi.org/10.1016/j.bioflm.2022.100071

Wang, G., Wang, F., Huang, Q., Li, Y., Liu, Y., & Wang, Y. (2015). Understanding transcription factor regulation by integrating gene expression and DNase I hypersensitive sites. *BioMed Research International, 2015*, 757530. https://doi.org/10.1155/2015/757530

Acknowledgements

This work was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant to J.L. Strap.